

# Instruction for CARBAMAZEPINE ™

Please ensure you have the correct prescription and order form in front of you. There are several different order forms for Antidepressants (24), Anti-psychotics (23), Sedatives (19), Anti-epileptics (7), Hormones (1) & Analgetics (4).

## PLEASE READ CAREFULLY BEFORE FILLING IN THE PRESCRIPTION AND ORDER FORM

1. A sudden termination of treatment with carbamazepine or a sudden reduction in dosage can lead to significant withdrawal symptoms. To avoid this, the daily dose should be reduced in small steps gradually over a period of time. This can be done using tapering strips™.
2. One or more tapering strips™ can be used to gradually reduce the dose of carbamazepine to a desired lower dose or to zero. If more than one strip is needed the final dose of the previous strip is always equal to the starting dose of the following strip.
3. The treating physician should plan the tapering schedule together with the patient. It is not possible to predict the perfect schedule for each patient. Some patients are far more sensitive to withdrawal symptoms than others, and some patients cope with symptoms better than others. The risk of withdrawal symptoms and their intensity can be reduced by choosing a more gradual tapering schedule. There are two options when choosing a tapering schedule:
  - 1) **Variable** - Choose a tapering schedule for the first 28 days only, then use the patient's experiences and feedback to inform the remainder of the tapering trajectory. To do this only the **black** part of table 4 on page 4 has to be filled in. The experiences of the patient during the first weeks (withdrawal symptoms or not) can then be used to choose the tapering trajectories for the following strip(s) by filling in in the **red** part of the table 5 on page 5.
  - 2) **Fixed** - Choose a tapering schedule for the entire period of the dosage reduction. To do this, fill in both the **black** and (if applicable) the **red** part of table 4 on page 4.
4. Carbamazepine tapering strips™ are only available on prescription. The signed prescription and order form is seen as a valid prescription. After filling in the form, the treating physician should terminate any current prescription for carbamazepine at the local pharmacy.
5. In most countries, the costs of the medication in the tapering strips™ are covered by health insurance. Tapering and stabilisation strips contain medication for a fixed period of 28 days, all strips have the same price. Tapering strip™: € 77.-; Stabilization strip € 38.50 for a period of 28 days; Prices are subject to change, so please check before ordering.
6. Once the prescription and order form has been received by the Regenboog Apotheek, the tapering strip™ will be shipped to the patient's home address (Europe, North America and Australia) within a week.
7. A tapering strip™ contains 28 numbered pouches, with each pouch containing several tablets. The number of pills may vary per day. The total dose of the medication reduces over time as the patient takes the pills in each daily pouch.
8. The patient starts their tapering schedule on a Sunday by taking the pills in the first pouch, numbered 028. **By counting down**, the patient finishes four weeks later on a Saturday, by taking the pills in the last pouch, numbered 001. By doing this, the patient can easily keep track of how many doses and days are left.
9. Each pouch of the strip is printed with the sequence number, the colour and strength of each tablet, the total number of tablets and the name of the pharmacy.
10. **STABILIZATION**: A stabilization strip can keep the patient on a fixed dose for a desired period of time. Such stabilization may be necessary if tapering appears to be going too fast for the patient, or the patient does not wish to reduce the dose to zero, but remain at an optimal lower dose. The stabilization strip can be produced at almost any desired dose rounded off to 5 mg carbamazepine.
11. In some cases, the patient is currently using a dose that is higher or different from the starting dose of the standard available tapering strips (see table 4 on page 4). These patients can also reduce their dose gradually by using **customized** tapering strips™ for the first part of the taper (see table 4 on page 4).

12. For carbamazepine, the following tapering strips™ are standard available:

Startdose carbamazepine	Enddose carbamazepine	Tapering period	Name of the strip	step dose reduction	average dose reduction	average dose reduction
400 mg/day	300 mg/day	28 days	CRBM 400-300/28days	5 mg	3,57 mg/day	6,3%/week
400 mg/day	200 mg/day	28 days	CRBM 400-200/28days	5 /10 mg	7,14 mg/day	12,5%/week
300 mg/day	200 mg/day	28 days	CRBM 300-200/28days	5 mg	3,57 mg/day	8,4%/week
300 mg/day	100 mg/day	28 days	CRBM 300-100/28days	5 /10 mg	7,14 mg/day	16,7%/week
200 mg/day	100 mg/day	28 days	CRBM 200-100/28days	5 mg	3,57 mg/day	12,5%/week
100 mg/day	50 mg/day	28 days	CRBM 100-50/28days	5 mg	1,79 mg/day	12,5%/week
100 mg/day	zero	28 days	CRBM 100-0/28days	5 mg	3,57 mg/day	25%/week
50 mg/day	20 mg/day	28 days	CRBM 50-20/28days	5 mg	1,07 mg/day	15%/week
50 mg/day	zero	28 days	CRBM 50-0/28days	5 mg	1,79 mg/day	25%/week
20 mg/day	zero	28 days	CRBM 20-0/28days	5 mg	0,71 mg/day	25%/week
stable	. . . mg/day	28 days	CRBM stable	0 mg	no reduction	0%/week

The table below shows how many mg of carbamazepine each daily pouch contains.

Schedule: carbamazepine		400 - 300	400 - 200	300 - 200	300 - 100	200 - 100	100 - 50	100 - 0	50 - 20	50 - 0	20 - 0	10 - 10
Sachet No	Intake day	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg
sachet 28	sunday	400	400	300	300	200	100	100	50	50	20	10
sachet 27	monday	395	395	295	295	195	100	95	50	50	20	10
sachet 26	tuesday	395	380	295	285	195	95	95	50	45	20	10
sachet 25	wednesday	390	380	290	280	190	95	90	45	45	20	10
sachet 24	thursday	380	370	285	270	185	95	85	45	45	15	10
sachet 23	friday	380	365	280	265	180	90	80	45	40	15	10
sachet 22	saturday	380	355	280	255	180	90	80	45	40	15	10
sachet 21	sunday	375	350	275	250	175	85	75	40	40	15	10
sachet 20	monday	370	340	270	240	170	85	70	40	35	15	10
sachet 19	tuesday	365	335	265	235	165	85	70	40	35	15	10
sachet 18	wednesday	365	325	265	225	165	80	65	40	30	15	10
sachet 17	thursday	360	320	260	220	160	80	60	40	30	10	10
sachet 16	friday	355	310	255	210	155	80	55	35	30	10	10
sachet 15	saturday	350	305	250	205	150	75	55	35	25	10	10
sachet 14	sunday	350	295	250	195	150	75	50	35	25	10	10
sachet 13	monday	345	290	245	190	145	70	45	35	25	10	10
sachet 12	tuesday	340	280	240	180	140	70	45	30	20	10	10
sachet 11	wednesday	335	275	235	175	135	70	40	30	20	10	10
sachet 10	thursday	335	265	235	165	135	65	35	30	20	5	10
sachet 9	friday	330	260	230	160	130	65	30	30	15	5	10
sachet 8	saturday	325	250	225	150	125	65	30	30	15	5	10
sachet 7	sunday	320	245	220	145	120	60	25	25	15	5	10
sachet 6	monday	320	235	220	135	120	60	20	25	10	5	10
sachet 5	tuesday	315	230	215	130	115	55	20	25	10	5	10
sachet 4	wednesday	310	220	210	120	110	55	15	25	5	5	10
sachet 3	thursday	305	215	205	115	105	55	10	20	5	5	10
sachet 2	friday	305	205	205	105	105	50	5	20	5	5	10
sachet 1	saturday	300	200	200	100	100	50	5	20	5	5	10

**IMPORTANT NOTICE: Tapering strips™ are not intended to replace medical care provided by a physician. Careful counselling and monitoring of the patient remains essential during and after treatment with tapering strips.**

Please note: Treatment with the tapering strip™ supports the tapering of medication, reducing withdrawal effects to a minimum. If withdrawal effects do occur, they will be not as intensive or prolonged as during a traditional tapering approach. This will make it easier for the treating physician to recognize a relapse (**such as a** return of depression or anxiety) and to distinguish between relapse or withdrawal effects in good time.

There is a noticeable difference in timing between the occurrence of withdrawal symptoms and relapse. Withdrawal symptoms usually occur shortly or immediately after the start of tapering. Relapse usually occurs at a later stage, often after tapering with carbamazepine has been completed. While withdrawal symptoms become less severe and disappear over time, symptoms as a result of relapse tend to remain and may even get worse.

Should you have any questions or if you require more information, please contact Paul Harder, pharmacist at the Regenboog Apotheek at [pharder@regenboogapotheek.nl](mailto:pharder@regenboogapotheek.nl) or + 31-(0)6-25072020.

## Instructions for the prescription and order form for tapering strips™

- MEDICAL INDICATION: please, mark all the indications applicable to the patient:
  - preventive** reducing the risk of withdrawal symptoms or relapse effects caused by diminishing the dose too quickly.
  - reduction anxiety** patient is excessively anxious about tapering, for example as a result of comorbid anxiety disorder.
  - failed attempt** previous attempts to stop or reduce the dose have failed.
  - distinction** distinguishing between withdrawal symptoms and possible relapse.
  - slow metabolizer** the lowest dose causes a high plasma concentration, increasing the risk of withdrawal effects.
  - high dose** the patient has used the medication for a period of 6 months at a dose at least 150% of the defined daily dose.
- Fill in the patient's details.
- Fill in the details of the treating physician.
- Fill in the table. Below you can see four examples showing the different ways of filling in the table:

- a. **Total reduction**, using **3** tapering strips™: mark **3** boxes  
(schedule 3 x 28 days = 84 days)

starting dose EXAMPLE	taper duration	STRIP 1 28 days	STRIP 2 28 days	STRIP 3 28 days	STRIP 4 28 days
30 mg/day	112 days	<input type="checkbox"/> EXMPL 30-10	<input type="checkbox"/> EXMPL 10-5	<input type="checkbox"/> EXMPL 5-2	<input type="checkbox"/> EXMPL 2-0
	84 days	<input checked="" type="checkbox"/> EXMPL 30-10	<input checked="" type="checkbox"/> EXMPL 10-5	<input checked="" type="checkbox"/> EXMPL 5-0	
	56 days	<input type="checkbox"/> EXMPL 30-10	<input type="checkbox"/> EXMPL 10-0		

- b. **Dose reduction** using **2** tapering strips™: mark **2** boxes  
(schedule 2 x 28 days = 56 days)

starting dose EXAMPLE	taper duration	STRIP 1 28 days	STRIP 2 28 days	STRIP 3 28 days	STRIP 4 28 days
30 mg/day	112 days	<input type="checkbox"/> EXMPL 30-10	<input type="checkbox"/> EXMPL 10-5	<input type="checkbox"/> EXMPL 5-2	<input type="checkbox"/> EXMPL 2-0
	84 days	<input checked="" type="checkbox"/> EXMPL 30-10	<input checked="" type="checkbox"/> EXMPL 10-5	<input type="checkbox"/> EXMPL 5-0	
	56 days	<input type="checkbox"/> EXMPL 30-10	<input type="checkbox"/> EXMPL 10-0		

- c. **Customized schedule** please mark each box, and fill in the desired start and end dose for each strip.  
Total duration = number of strips x 28 days.

<b>CUSTOMIZED</b> 2 x 28 days	<input checked="" type="checkbox"/> EXMPL from <b>60 - 30</b> mg	<input checked="" type="checkbox"/> EXMPL from <b>30 - 10</b> mg
-------------------------------	--	--

- d. **Stabilization** to a steady dose: mark the box, and fill in daily dose and number of repeats  
(in this example: 17 mg/day and 2x repeats=> schedule (1 + 2) x 28 days = 84 days)

<b>STABILIZATION</b> 28 days	<input checked="" type="checkbox"/> EXMPL stab.	<b>17</b> mg/day	<b>2</b> repeats
------------------------------	---	------------------	------------------

- Sign the prescription.
- Fax the prescription to the Regenboog Apotheek: **+31-(0)85-2736129**

**NOTE: Please only fax the order and prescription form.**

Do not fax the instruction sheets.

# Order and prescription form

**R/ Prescription = order form for CABAMAZEPINE tapering strips™** ZI-number: 14729709

- 1a.** Medical indication:  preventive  reduction anxiety  failed attempt  distinction  slow metabolizer  high dose  
**1b.** Duration of carbamazepine use:  <1 year  1-2 years  2-5 years  5-10 years  >10 years

**2.** Full name of patient: .....

Date of birth: . . . - . . . - . . . . . gender:  M /  F

Address: .....

City, postal code and country: .....

E-mail address: ..... Telephone no: .....

**3.** Name of treating physician: .....

Medical registration number: .....

Address: .....

Postal code and country: .....

E-mail address: .....

Telephone no: .....

**4. Please, fill in the table, following the instructions above.**

Startdose CARBAMAZEPINE	Tapering period	STRIP 1 28 days	STRIP 2 28 days	STRIP 3 28 days	STRIP 4 28 days	STRIP 5 28 days
400 mg/day	140 days	<input type="checkbox"/> CRBM 400-300	<input type="checkbox"/> CRBM 300-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0
	112 days	<input type="checkbox"/> CRBM 400-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0	
	84 days	<input type="checkbox"/> CRBM 400-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-0		
300 mg/day	112 days	<input type="checkbox"/> CRBM 300-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0	
	84 days	<input type="checkbox"/> CRBM 300-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-0		
	56 days	<input type="checkbox"/> CRBM 300-100	<input type="checkbox"/> CRBM 100-0			
200 mg/day	84 days	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0		
	56 days	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-0			
100 mg/day	56 days	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0			
	28 days	<input type="checkbox"/> CRBM 100-0				
50 mg/day	56 days	<input type="checkbox"/> CRBM 50-20	<input type="checkbox"/> CRBM 20-0			
	28 days	<input type="checkbox"/> CRBM 50-0				
20 mg/day	28 days	<input type="checkbox"/> CRBM 20-0				
<b>customized</b>	. X 28 days	<input type="checkbox"/> CRBM . . . . .	<input type="checkbox"/> CRBM . . . . .	<input type="checkbox"/> CRBM . . . . .	<input type="checkbox"/> CRBM . . . . .	<input type="checkbox"/> CRBM . . . . .
	set-up 1 X . . days	<input type="checkbox"/> CRBM . . . . .				
<b>stable</b>	28 days	<input type="checkbox"/> CRBM stable	. . . mg/day	. . repeats	dose round off to 5 mg	please send immediately

**COSTS** Tapering and stabilisation strips contain medication for a fixed period of 28 days, all strips have the same price.  
 Tapering strip™: € 77.-; Stabilization strip € 38.50 for a period of 28 days; Prices are subject to change.

**5.** Date: ..... Stamp of treating physician

Signature of treating physician:

Shipping address: Regenboog Apotheek, Antwoordnummer 16500, 4840 WJ BAVEL, The Netherlands

# Order and prescription form

**R/ Prescription = order form for CABAMAZEPINE tapering strips™**

ZI-number: 14729709

**1. Continuation Tapering**

**2. Full name of patient:** .....

Date of birth: . . . - . . . - . . . . . gender:  M /  F

Address: .....

City, postal code and country: .....

E-mail address: ..... Telephone no: .....

**3. Name of treating physician:** .....

Medical registration number: .....

Address: .....

Postal code and country: .....

E-mail address: .....

Telephone no: .....

**5. Please, fill in the table, following the instructions above.**

Startdose CARBAMAZEPINE	Tapering period	STRIP 1 28 days	STRIP 2 28 days	STRIP 3 28 days	STRIP 4 28 days	STRIP 5 28 days
400 mg/day	140 days	<input type="checkbox"/> CRBM 400-300	<input type="checkbox"/> CRBM 300-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0
	112 days	<input type="checkbox"/> CRBM 400-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0	
	84 days	<input type="checkbox"/> CRBM 400-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-0		
300 mg/day	112 days	<input type="checkbox"/> CRBM 300-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0	
	84 days	<input type="checkbox"/> CRBM 300-200	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-0		
	56 days	<input type="checkbox"/> CRBM 300-100	<input type="checkbox"/> CRBM 100-0			
200 mg/day	84 days	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0		
	56 days	<input type="checkbox"/> CRBM 200-100	<input type="checkbox"/> CRBM 100-0			
100 mg/day	56 days	<input type="checkbox"/> CRBM 100-50	<input type="checkbox"/> CRBM 50-0			
	28 days	<input type="checkbox"/> CRBM 100-0				
50 mg/day	56 days	<input type="checkbox"/> CRBM 50-20	<input type="checkbox"/> CRBM 20-0			
	28 days	<input type="checkbox"/> CRBM 50-0				
20 mg/day	28 days	<input type="checkbox"/> CRBM 20-0				
<b>customized</b>	. X 28 days	<input type="checkbox"/> CRBM . . . . .	<input type="checkbox"/> CRBM . . . . .	<input type="checkbox"/> CRBM . . . . .	<input type="checkbox"/> CRBM . . . . .	<input type="checkbox"/> CRBM . . . . .
set-up	1 X . . days	<input type="checkbox"/> CRBM . . . . .				
<b>stable</b>	28 days	<input type="checkbox"/> CRBM stable	. . . mg/day	. . repeats	dose round off to 5 mg	please send immediately

**COSTS** Tapering and stabilisation strips contain medication for a fixed period of 28 days, all strips have the same price. Tapering strip™: € 77.-; Stabilization strip € 38.50 for a period of 28 days; Prices are subject to change.

**6. Date:**

Stamp of treating physician

Signature of treating physician:

Shipping address: Regenboog Apotheek, Antwoordnummer 16500, 4840 WJ BAVEL, The Netherlands

SUMMARY, 24 May 2018

## **Antidepressant tapering strips to help people come off medication more safely**

Peter Groot and Jim van Os

Antidepressant users who have come to the end of treatment and wish to stop can suffer withdrawal symptoms, even when following their doctor's advice. In some cases, these withdrawal symptoms can be so severe that they prevent the user from stopping their medication. This can often be problematic for the patient and can result in unnecessary prescriptions, additional cost for health services and potential long-term health issues.

An observational study published in the journal *Psychosis* shows that antidepressant withdrawal symptoms can be minimised or avoided by using prepackaged medication in tapering strips. Tapering medication makes safe withdrawal possible by utilising pills of small dosages to make much smaller dosage reductions than can be achieved using standard registered dosages. This approach makes tapering easy for the antidepressant user and the doctor. The study shows that more patients are successful using tapering strips when compared to those who had tried to reduce using other methods and allows patients who have previously tried and failed to withdraw the opportunity to succeed in coming off their antidepressant medication.

### **What is tapering medication?**

Tapering medication is prepackaged in tapering strips which make it easy for the patient to reduce without needing to follow complicated instructions. Furthermore the strips make it easy for doctors to prescribe personalised tapering schedules for their patients. A tapering strip consists of 28 sachets on a roll, with each sachet providing the medication for a single day. The daily dosage in each sachet is a little less or equal to that of the previous day. A doctor, psychiatrist or therapist can prescribe tapering schedules tailored to and in consultation with the patient in a shared decision-making approach. Using currently available standard registered dosages, gradual tapering is often difficult and, in some cases, impossible to achieve. Due to these limitations, current tapering guidelines are also difficult to follow for both the prescriber and the patient.

### **About the study**

The study reviewed 1121 questionnaires completed by patients who wanted to withdraw from their antidepressant medications. The majority (75%,  $n = 895$ ) wanted to taper off completely. Of these, 62% ( $n = 692$ ) had previously made one or more tapering attempts. Almost everyone (97%) had suffered withdrawal symptoms during these earlier attempts, which were rated as very severe by 49% (7 on a scale of 1-7). The duration of antidepressant use varied from less than a year to more than 30 years (median: 2-5 years).

### **A comparison of withdrawal with or without tapering medication**

All patients were asked about their tapering experience, using a scale from 1-7, where 1 = very good and 7 = very bad. In addition, they were asked if they had experienced withdrawal symptoms on a 1-7 scale, where 1 = not at all and 7 = very much. Patients were also asked to rate their previous attempts to withdraw, therefore making a comparison possible between tapering medication and other methods.

### **Study findings**

The main results of the study are that, of the 895 people who used tapering strips to taper the antidepressant they used:

- 71% ( $n = 636$ ) succeeded in a median time of 56 days (2 tapering strips).

- 21% (n = 192) did not complete the taper, 8% (n = 67) because they were still tapering, 4% (n = 39) due to withdrawal symptoms and 6% (n = 53) due to relapse (defined as the recurrence of complaints such as anxiety or depression).

The results of this study show that, for patients who had previously tried and failed to withdraw, the use of tapering medication in tapering strips could enable them to withdraw successfully.

### **Tapering medication enables shared decision making**

For patients who wish to withdraw, attempting to taper using standard registered doses increases the risk of withdrawal symptoms, leading to an increased risk of withdrawal failure and unnecessary long-term antidepressant use. Allowing patients and doctors to work together to define personalised tapering schedules has been shown to reduce withdrawal symptoms and increase the chances of success. By doing this, doctors can focus on patient recovery and more easily distinguish between withdrawal symptoms and relapse, which is important in avoiding the unnecessary prescribing of antidepressant medications.

### **Applying tapering principles to other medications**

The results of this study are relevant for people using other psychotropic medications such as antipsychotics or benzodiazepines. This is important for people with psychosis who will often be taking multiple medications. The use of tapering medication in tapering strips offers the opportunity to make safe, gradual dosage adjustments while minimising unpleasant withdrawal symptoms even in those patients who are taking multiple medications. The use of tapering strips enables a level of flexibility and personalisation that is simply not achievable using conventional standard registered dosages.

---

Information about taperingstrips can be found on the User Research Center of Maastricht University website [www.taperingstrip.org](http://www.taperingstrip.org)

Dr. Peter C. Groot is a researcher and expert by experience at the User Research Center of Maastricht University and volunteer at the Cinderella Therapeutics Foundation. Prof. Jim van Os is Chairman of the Brain Division at the Utrecht University Medical Center.

Email: [p.c.groot@maastrichtuniversity.nl](mailto:p.c.groot@maastrichtuniversity.nl), [j.i.vanos-2@umcutrecht.nl](mailto:j.i.vanos-2@umcutrecht.nl).

---